

# PROVIDER TIPS

## Pediatric HEDIS Measures: At A Glance

Measures	Coding Tips	Recommendations
<p><b>Child and Adolescent Well-Care Visits (WCV)</b> 3-21 years of age who have had at least one comprehensive well-care visit with a PCP or an OB/GYN practitioner during the measurement year.</p>	<p><b>ICD10 Codes</b> Z00.121-Z00.129, Z00.2, Z00.3, Z02.5, Z76.1-Z76.2.</p> <p><b>CPT Codes</b> 99381-99385, 99391-99395, 99461</p> <p><b>HCPCS Codes</b> G90438, G0439, S0302</p>	<ul style="list-style-type: none"> <li>Take advantage of each office visit (including sick visits) to provide well-care services and immunizations. To bill both components on the same day, add Modifier -25 to the sick visit and billing for the appropriate preventative visit.</li> </ul> <p><b>The comprehensive well-care must include:</b></p> <ul style="list-style-type: none"> <li>✓ Health History: assessment of Member's history of disease or illness and family health history.</li> <li>✓ Physical Development History: assessment of specific age-appropriate physical development milestones.</li> <li>✓ Mental Development History: assessment of specific age-appropriate mental development milestones.</li> <li>✓ Physical exam.</li> <li>✓ Health Education/Anticipatory Guidance: guidance given in anticipation of emerging issues that a child/family may face.</li> </ul> <ul style="list-style-type: none"> <li>Use standardized templates in charts and in EMRs that allow check boxes documentation of counseling.</li> <li>Schedule next visit at end of each appointment.</li> </ul>
<p><b>Well-Child Visits in the First 30 Months of Life (W30)</b></p> <p><b>1. Well-Child Visits in the First 15 Months</b> Children who turned 15 months old during the measurement year: six or more well-child visits.</p> <p><b>2. Well-Child Visits for Age 15-30 Months</b> Children who turned 30 months old during the measurement year: two or more well-child visits.</p>	<p><b>ICD10 Codes</b> Z00.110, Z00.111, Z00.121, Z00.129, Z76.1-Z76.2</p> <p><b>CPT Codes</b> 99381-99385, 99391-99395, 99461</p> <p><b>HCPCS Codes</b> G90438, G0439, S0302</p>	<ul style="list-style-type: none"> <li>Use standardized templates in charts and in EMRs that allow check boxes documentation of counseling.</li> <li>Schedule next visit at end of each appointment.</li> </ul>
<p><b>Weight Assessment and Counseling for Nutrition and Physical Activity</b> Age 3-17 years</p>	<p><b>BMI Percentile</b> ICD-10*: Z68.51-Z68.54</p> <p><b>Counseling for Nutrition</b> ICD-10*: Z71.3 CPT: 97802-97804 HCPCS: G0270, G0271, G0447</p> <p><b>Counseling for Physical Activity</b> ICD 10: Z02.05, Z71.82 HCPCS: G0447</p>	<ul style="list-style-type: none"> <li>Take advantage of each office visit (including sick visits) to counsel for nutrition and physical activity. <b>Discuss and document nutrition and physical activity during at least one office visit annually.</b></li> <li>At each office visit, document BMI percentile (to include height, weight and BMI percentile).</li> <li>Nutrition Counseling: discuss current nutrition behaviors (e.g., appetite or meal patterns, eating and dieting habits).</li> <li>Physical Activity: discuss current physical activity behaviors, exercise routine, sports activities, sports physical, weight or obesity counseling.</li> </ul>

Measures	Coding Tips	Recommendations
<p><b>URI - Appropriate Treatment for Children with Upper Respiratory Infection</b></p> <p>Age 3 months to 18 years that were given only a diagnosis of URI and were NOT dispensed an antibiotic prescription.</p>	<p><b>Pharyngitis ICD10 Codes</b> J02.0, J02.8–J03.01, J03.80–J03.81, J03.90–J03.91</p> <p><b>Group A Strep Tests CPT Codes</b> 87070, 87071, 87081, 87430, 87650–87652, 87880</p>	<ul style="list-style-type: none"> <li>• Test all children for group A strep before prescribing an antibiotic for a diagnosis of pharyngitis only.</li> <li>• Document and submit claims for all appropriate diagnoses established at the visit.</li> <li>• Submit claim for in-office rapid strep test.</li> <li>• Do not prescribe antibiotics for URI diagnosis only.</li> </ul>
<p><b>Childhood Immunizations with Lead Screening</b></p> <p>Age 2 years who have had the following vaccines on or before their second birthday:</p> <ul style="list-style-type: none"> <li>• 4 DTaP (diphtheria, tetanus and acellular pertussis)</li> <li>• 3 IPV (polio)</li> <li>• 1 MMR (measles, mumps, rubella)</li> <li>• 3 HiB (H influenza type B)</li> <li>• 3 Hep B (hepatitis B)</li> <li>• 1 VZV (chicken pox)</li> <li>• 4 PCV (pneumococcal conjugate)</li> <li>• 1 Hep A (hepatitis A)</li> <li>• 2 or 3 RV (rotavirus)</li> <li>• 2 Flu (Influenza) age: 0–2yrs</li> </ul> <p>Lead screening: on or before age two, by second birthday.</p>	<p><b>DTaP-HiB-IPV</b> 90698</p> <p><b>DTaP-HepB-IPV</b> 90723</p> <p><b>IPV*</b> 90713</p> <p><b>MMR*</b> 90707</p> <p><b>MMRV</b> 90710</p> <p><b>HiB*</b> 90647 or 90648 with (90460/90641 or 90471/90472)</p> <p><b>HepB*</b> 90473, 90744 or 90746 with (90460/90461 or 90471/90472)</p> <p><b>VZV</b> 90716</p> <p><b>PCV13</b> 90670 with (90460/90461 or 90471/90472)</p> <p><b>Influenza</b> 90630, 90657, 90661, 90673, 90685 with (90460/ 90461 or 90473/90474)</p> <p><b>Rotavirus</b> Two dose: 90681 with (90460/ 90461 or 90473/90474) Three dose: 90680 with (90460/ 90461 or 90473/90474)</p> <p><b>Lead CPT Code</b> 83655</p> <p><b>ICD10 Code</b> Z23</p> <p><b>CPT Code</b> 3E0234Z</p>	<ul style="list-style-type: none"> <li>• Review immunization record at each visit and administer age-appropriate vaccines.</li> <li>• Establish a reminder system for patients.</li> <li>• Take advantage of every office visit (including sick visits) to catch up on missing vaccines.</li> <li>• Use the State Immunization Registry.</li> <li>• Lead screening test should be completed on all children before their second birthday. A lead risk questionnaire does not count– the screening must be a capillary or venous blood lead test.</li> </ul>

Measures	Coding Tips	Recommendations
<p><b>IMA - Immunizations in Adolescents</b></p> <p>Members who turned 13 years of age in the measurement year and received by age 13.</p> <ul style="list-style-type: none"> <li>• <b>Tdap Vaccine:</b> one dose between 10th and 13th birthday.</li> <li>• <b>Meningococcal Conjugate Vaccine:</b> one dose of meningococcal Y vaccine between 11th and 13th birthday.</li> <li>• <b>HPV Vaccine:</b> either <ul style="list-style-type: none"> <li>- two doses of HPV vaccine between 9th and 13th birthday with at least 146 days between doses, or</li> <li>- three doses with different dates of service between 9th and 13th birthday.</li> </ul> </li> </ul>	<p><b>Tdap</b></p> <p><b>CPT Code</b> 90714, 90715, <b>CVX Code</b> 90718 115</p> <p><b>Meningococcal</b></p> <p><b>CPT Code</b> 90734 <b>CVX Codes</b> 108, 114, 136, 147, 167</p> <p><b>HPV</b></p> <p><b>CPT Codes</b> 90649, 90650, 90651 <b>CVX Codes</b> 62, 118, 137, 165</p>	<ul style="list-style-type: none"> <li>• Review immunization record at each visit and administer age-appropriate vaccines.</li> <li>• Establish a reminder system for patients.</li> <li>• Take advantage of every office visit (including sick visits) to catch up on missing vaccines.</li> <li>• Use the State Immunization Registry.</li> </ul>